


1/6

Library	Sequence	Incidence
Ph.D.-12	HPLKQYWWRPSI-----	22/26
Ph.D.-12	---PIWWKHSGGPIL-	1/26
Ph.D.-12	---YWWRDAPVSQGR	1/26
Ph.D.-12	SYPTDKWWIKPG-----	1/26
Ph.D.-7	---VQWWKRPT-----	7/15
Ph.D.-7	---NWWKPLP-----	1/15
Ph.D.-7	---GKWWVFD-----	1/15
Ph.D.-7	-VPTKPKWW-----	1/15
Ph.D.-C7C	----PWWKTSK-----	6/15
Ph.D.-C7C	----PWWKASS-----	1/15
Ph.D.-C7C	----TPTWWRT-----	1/15
Ph.D.-C7C	----APTWWKS-----	1/15
Ph.D.-C7C	----WWTSASR-----	1/15
Ph.D.-C7C	----SARWWQP-----	1/15

FIGURE 1

BEST AVAILABLE COPY

2/6



BEST AVAILABLE COPY

FIGURE 2

3/6




FIGURE 3

A/6

BEST AVAILABLE COPY




FIGURE 4

BEST AVAILABLE COPY





FIGURE 5

6/6

Suggested Amino Acid Substitutions.
solvent exposed (SEA^a>30 Å²) / interior (SEA^a<10 Å²)



Amino acids connected by a solid line can be substituted with 95% confidence
(D. Bordo and P. Argos, J. Mol. Biol., 217(1991)721-729)

^aSEA=solvent exposed area

Figure 6

00200400 - 060100

BEST AVAILABLE COPY